astro8405

An Introduction to the Cosmic Microwave Background

Kaustuv Basu

kbasu@uni-bonn.de



eCampus | Lernplattform der Universität Bonn



astro8405: The Cosmic Microwave Background

Aktionen 🕶


This course intends to give you a modern and up-to-date introduction to the science and experimental techniques relating to the Cosmic Microwave Background. No prior knowledge of cosmology is necessary, your prerequisite are a basic understanding of electrodynamics and thermal physics and some familiarity with Python programming.

Lecture 9:

Polarization Anisotropies:

B-modes and the primordial gravitational waves

Recap: E- and B-modes of polarization



CMB has weak linear polarization (Thomson scattering does not create circular polarization). Polarization anisotropies are described by a vector field, whereas temperature anisotropies are represented by a scalar field.

The rotational transformation of Stokes Q and U parameters lead us to define two coordinate-free variables, E and B, using the properties of spin-weighted spherical harmonics.

$$T(\hat{\boldsymbol{n}}) = \sum_{lm} a_{T,lm} Y_{lm}(\hat{\boldsymbol{n}})$$
$$(Q + iU)(\hat{\boldsymbol{n}}) = \sum_{lm} a_{2,lm} \, _2Y_{lm}(\hat{\boldsymbol{n}})$$
$$(Q - iU)(\hat{\boldsymbol{n}}) = \sum_{lm} a_{-2,lm} \, _{-2}Y_{lm}(\hat{\boldsymbol{n}}).$$

$$\tilde{E}(\hat{\boldsymbol{n}}) = \sum_{lm} \left[\frac{(l+2)!}{(l-2)!} \right]^{1/2} a_{E,lm} Y_{lm}(\hat{\boldsymbol{n}})$$

$$\tilde{B}(\hat{\boldsymbol{n}}) = \sum_{lm} \left[\frac{(l+2)!}{(l-2)!} \right]^{1/2} a_{B,lm} Y_{lm}(\hat{\boldsymbol{n}})$$

$$a_{E,lm} = -(a_{2,lm} + a_{-2,lm})/2$$


 $a_{B,lm} = i(a_{2,lm} - a_{-2,lm})/2.$

Recap: E and B mode characteristics


Two flavours of CMB polarization:


E mode → even parity → scalar perturbations B mode → odd parity → tensor perturbations

- We break down the polarization field into two components which we call E and B modes. This is the spin-2 analog of the gradient/curl decomposition of a vector field.
- E modes are generated by density (scalar) perturbations via Thomson scattering.
- Additional vector modes are created by vortical motion of the matter at recombination – this is small
- B modes are generated by gravity waves (tensor perturbations) at last scattering, or by gravitational lensing (which transforms E modes into B modes along the line of sight to us) later on.









What causes the CMB quadrupole?

Two things:

"Normal" CDM: Density perturbations at z≈1100 lead to velocities that create local quadrupoles seen by scattering electrons.

=> E-mode polarization ("parity-even")

Gravity waves: create local quadrupoles seen by the scattering electrons.

=> B-mode polarization ("parity-odd")

The problem of understanding the polarization pattern of the CMB thus reduces to understanding the quadrupole temperature fluctuations at the *moment of last scattering*.


From gravity waves to B-mode polarization

Arguably the cleanest prediction of inflation is a spectrum of primordial gravitational waves. These are tensor perturbations to the spatial metric,

$$ds^2 = a^2 \sum_{i=1}^3 \sum_{j=1}^3 (\delta_{ij} + h_{ij}) dx^i dx^j$$

-> CURVED SPACE!

A plane gravitational wave causes a "quadrupolar stretching" of the space (tensor mode perturbations, as opposed to scalar modes from density).



From gravity waves to B-mode polarization

Arguably the cleanest prediction of inflation is a spectrum of primordial gravitational waves. These are tensor perturbations to the spatial metric,


$$ds^2 = a^2 \sum_{i=1}^3 \sum_{j=1}^3 (\delta_{ij} + h_{ij}) dx^i dx^j$$

"metric perturbation"

-> CURVED SPACE!


A plane gravitational wave causes a "quadrupolar stretching" of the space (tensor mode perturbations, as opposed to scalar modes from density).

This changes a circle of test particles into an ellipse, and the radiation acquires a m=2 quadrupole pattern \rightarrow primordial B-mode signal




Gravitational waves detected by LIGO → wavelength thousands of kilometers

Primordial gravitational waves produced by inflation → wavelength billions of light-years!





GW propagating in isotropic electro-magnetic fields




 D_{ij} : Tensor metric perturbation [=gravitational waves]


$$ds^2=a^2\sum_{i=1}^3\sum_{j=1}^3(\delta_{ij}+h_{ij})dx^idx^j$$
 (D_{ij} and h_{ij} are the same thing, just different notations!) 09: Polarization anisotropies (part 2) 9







Why the GW changes photon energy




The change in photon energy (or its wavelength) is proportional to the time derivative of the metric perturbation, i.e., the difference in the metric perturbation at different epochs. As a result of inflation, the gravitational waves are very large (many times the horizon scale), hence the difference of metric perturbations between two different cosmic times is also large, resulting in the quadrupole temperature anisotropy from gravitational waves.

Why inflation would produce GW?

(From Kamionkowski & Kovetz, ARAA 2016)


Black hole analogy



HEURISTIC UNDERSTANDING OF INFLATIONARY GRAVITATIONAL WAVES


Here we present in simple heuristic terms the origin of IGWs. Consider first a black hole. As shown in Figure 2, it has an event horizon, a spherical surface beyond which, according to (classical) general relativity, objects and information disappear without a trace. Hawking showed, however, that when quantum mechanics is taken into consideration, the horizon glows—it emits electromagnetic radiation (Hawking 1975). Hawking's argument also applies, however, to any radiation field with massless quanta, and so the black hole also radiates GWs. In an FRW Universe with an accelerated expansion, there is also a horizon, a spherical surface beyond which (according to general relativity) objects and information disappear. This time, though, the observer sees this spherical surface from the inside, rather than the outside. Just as was the case with the black hole horizon, this horizon also radiates GWs, according to quantum mechanics. These GWs are produced throughout inflation, and the expansion rate and thus horizon temperature are nearly constant during inflation. These GWs thus remain, after inflation, as a primordial-GW background with a nearly scale-invariant spectrum.

Gravity waves also create E-modes



We should note that the tensor perturbations, caused by gravitational waves, also generate E-mode polarization (as well as temperature anisotropies). In fact, E- and B-modes are created almost equally (see figure above). However, the E-modes created by density (scalar) perturbations at the last scattering are much stronger, so one can neglect the tensor-origin of E-modes.

Recap on inflation and the meaning of the tensor-to-scalar ratio, *r*



I. Scalar power spectrum

The scaler perturbations are Gaussian, so all information about them is contained in the two-point correlation function:

 $\langle \mathcal{R}(\mathbf{k})\mathcal{R}^*(\mathbf{k}')\rangle = \frac{P(k)}{(2\pi)^3}\delta(\mathbf{k} - \mathbf{k}'),$

The mean square value of the initial perturbation amplitude is

$$\langle \mathcal{R}^2(\mathbf{x}) \rangle = \langle \int e^{i\mathbf{k}\mathbf{x}} R(\mathbf{k}) d^3k \int e^{-i\mathbf{k}'\mathbf{x}} R^*(\mathbf{k}') d^3k' \rangle = \int d^3k \frac{P(k)}{(2\pi)^3} = \int_0^\infty \frac{dk}{k} \mathcal{P}(k),$$

Where $\mathcal{P}(k) = k^3 P(k)/(2\pi^2)$ is also called the power spectrum, and is approximated as follows:

$$\mathcal{P}_s(k) = A_s \left(rac{k}{k_*}
ight)^{n_s-1}$$

In 1960's, Zel'dovich and Harrison independently predicted the flat spectrum of perturbations ($n_s=1$). But we now know that the spectrum is *slightly red*. The WMAP5 values for a fixed $k_*=500$ Mpc⁻¹ are:

$$A_s = (2.46 \pm 0.09) \cdot 10^{-9},$$

 $n_s = 0.960 \pm 0.014.$

2. Tensor power spectrum & power ratio, r

Actually, the derivation of approximately flat power spectrum does not depend on whether we deal with scalar or tensor fields. **Inflation also generates tensor perturbations** (transverse traceless perturbations of spatial metric h_{ij} , i.e. gravitational waves).

We have the same picture for tensor perturbations: primordial perturbations are Gaussian random field with almost flat power spectrum. In this case we have

$$\mathcal{P}_T(k) = A_T \left(rac{k}{k_*}
ight)^{n_T}.$$

It is convenient to introduce the parameter $r = \mathcal{P}_T/\mathcal{P}_s$ which measures the ratio of tensor to scalar perturbations.

For some early inflation theories with power-law potentials, prediction was $r \sim 0.1 - 0.3$

→ these are now practically ruled out by Planck data


3. Inflation and the spectral index, n_s

Inflation occurs if the universe is filled with a scalar field ϕ , which has non-vanishing scalar potential $V(\phi)$. The homogeneous field ϕ then satisfies the equation

$$\ddot{\varphi} + 3H\dot{\varphi} = -\frac{dV}{d\varphi}. \qquad \qquad a(t) \propto \exp\left(\int H dt\right), \quad H \approx {\rm const.}$$

For a relatively flat potential (dV/d ϕ small), the acceleration term can be neglected. The Friedmann equation in this case is H² = $8\pi/3G$ V(ϕ). So if ϕ varies slowly, then V(ϕ) and thus H also varies slowly, and the parameters of inflation are almost time independent (slow-roll inflation).

Yet, the parameters are not *exactly* time-independent (inflation has to end!), so the predicted value of the spectral tilt ($n_s - 1$) is small but non-zero. It can be positive or negative, depending on the scalar potential V(ϕ). In particular, it is negative for the simplest power-law potentials like



$$V(\varphi) = \frac{m^2}{2} \varphi^2 \quad \text{or} \quad V(\varphi) = \frac{\lambda}{4} \varphi^4.$$

For the case of slow-roll inflation,

$$n_s - 1 = -3M_{\rm pl}^2 \left(\frac{V'}{V}\right)^2 + 2M_{\rm pl}^2 \frac{V''}{V}$$
 with $' \equiv \frac{\partial}{\partial \phi}$.

4. Possible shapes of inflation potential




Figure 1

Inflation postulates that at some point in the early history of the Universe, the cosmic energy density was dominated by the vacuum energy associated with the displacement of some scalar field ϕ (the inflaton) from the minimum of its potential. Shown here for illustration are two toy models for the inflaton potential: Panel a shows a quadratic potential, and panel b shows a hilltop potential. Adapted from Kamionkowski & Kosowsky (1999) with permission.

$$\epsilon = 3 \frac{\dot{\phi}^2/2}{V + \dot{\phi}^2/2} \simeq \frac{M_{\rm Pl}^2}{2} \left(\frac{V'}{V}\right)^2$$
 The number of *e-folding* during inflation is determined by the slope of the potential, characterized by the parameter $\epsilon \equiv -\dot{H}/H^2 < 1$

$$N(t) \equiv \ln \frac{a(t_{\rm end})}{a(t)} = \int_{t}^{t_{\rm end}} H \, dt = -\frac{1}{2m_{\rm Pl}^{2}} \int_{\phi_{t}}^{\phi_{\rm end}} \frac{H}{H'} \, d\phi = \int_{\phi_{\rm end}}^{\phi_{t}} \frac{d\phi}{M_{\rm Pl}} \frac{1}{\sqrt{2\epsilon(\phi)}}.$$

5. Predictions for single-field, slow-roll inflation

Slow-Roll inflation requires the acceleration term for the potential is zero, and the shape of the potential is parametrized by the "slow-roll parameters"

$$\ddot{\varphi} + 3H\dot{\varphi} = -V'(\varphi)$$

$$H^{2} = \frac{V(\varphi)}{3M_{\rm Pl}^{2}}$$
$$3H\dot{\varphi} = -V'(\varphi)$$

$$H^{2} = \frac{V(\varphi)}{3M_{\rm Pl}^{2}} \qquad \qquad \varepsilon(\varphi) \equiv \frac{1}{2}M_{\rm Pl}^{2} \left(\frac{V'}{V}\right)^{2}$$
$$3H\dot{\varphi} = -V'(\varphi) \qquad \qquad \eta(\varphi) \equiv M_{\rm Pl}^{2} \frac{V''}{V}$$

For all physical, slow-roll inflation models, ϵ , $\eta << 1$

For scalar perturbations:

$$\Delta_{\mathcal{R}}^2(k) \equiv A_s \left(\frac{k}{k_{\star}}\right)^{n_s - 1} \qquad \boxed{n_s - 1 = -2\varepsilon - \eta}$$

$$n_s - 1 = -2\varepsilon - \eta$$

For tensor perturbations:

s:
$$r \equiv \frac{\Delta_t^2}{\Delta_R^2} = 16\epsilon$$

$$\Delta_t^2(k) \equiv A_t \left(\frac{k}{k_{\star}}\right)^{n_t}$$

$$r \equiv \frac{\Delta_t^2}{\Delta_R^2} = 16\epsilon$$

$$n_t = \frac{d \ln \Delta_t^2(k)}{d \ln k} = -2\epsilon$$

$$n_t = -2\epsilon$$

$$r \equiv \frac{\Delta_t^2}{\Delta_R^2} = 16\epsilon$$

$$n_t = \frac{\mathrm{d} \ln \Delta_t^2(k)}{\mathrm{d} \ln k} = -2\epsilon$$


$$r = 16\varepsilon$$
$$n_t = -2\varepsilon$$


The tensor amplitude is therefore a direct measure of the expansion rate H during inflation. Knowing this, one can completely determine the energetics of inflation!

- Single-field slow-roll inflation looks remarkably good:
 - Super-horizon fluctuation
 - Adiabaticity
 - Gaussianity
 - n_s<1


But we want a direct confirmation of inflation and probe its energy scale:

Gravitational waves!





Measurements of E, B power from ground




BICEP2 result in 2014

arXiv.org > astro-ph > arXiv:1403.3985

Astrophysics > Cosmology and Nongalactic Astrophysics

[Submitted on 17 Mar 2014 (v1), last revised 23 Jun 2014 (this version, v3)]

BICEP2 I: Detection Of B-mode Polarization at Degree Angular Scales



24


BICEP2 observation of the CMB polarization

0 h**CMB** 18 h


 μK_{CMB}

15

- Small telescope at South Pole
- 512 bolometers at 150 GHz
- Observed 380 square degrees for three years (2010 2012)
- Previous BICEP1 at 100 and 150 GHz (2006-2008)
- Current: Keck Array = $5 \times BICEP2$ at 150 GHz (2011 2013) and additional detectors at 100 and 220 GHz (2014 onwards)




BICEP2 E- and B-mode CMB maps




Detection of $\emph{B}\text{-}\text{Mode}$ Polarization at Degree Angular Scales by BICEP2

P. A. R. Ade et al. (BICEP2 Collaboration)
Phys. Rev. Lett. **112**, 241101 – Published 19 June 2014

BICEP2 scaler-to-tensor ratio




Then... proved wrong by *Planck*!



Planck's view of the BICEP2 field


colors → dust intensity

"engravings" → magnetic field




Credit: ESA/Planck collaboration

Planck results on the B-mode



Dust polarization map (Planck collaboration 2015)

Lensing B-mode (Planck collaboration 2015)



Discovery of Big Bang's Gravitational Waves Goes Bust, Due to Dust

A leaked analysis suggests that findings of gravitational waves from the big bang were illusory.

nature

Explore content > About the journal > Publish with us >

nature > news > article

Published: 30 January 2015

Gravitational waves discovery now officially dead

Ron Cowen


Nature (2015) | Cite this article

775 Accesses | 4 Citations | 677 Altmetric | Metrics


Combined data from South Pole experiment BICEP2 and Planck probe point to Galactic dust as confounding signal.

Cosmic inflation: Dust finally settles on BICEP2 results


by Amina Khan, Los Angeles Times




Planck 2015 + BICEP



BICEP2 + Planck joint analysis (2015)




95% upper limit: r < 0.08




32


Latest constraints (late 2021)


BICEP / Keck XIII: Improved Constraints on Primordial Gravitational Waves using Planck, WMAP, and BICEP/Keck Observations through the 2018 Observing Season



 n_{s}







SO will start observing very soon..







The Eternal Sky: A Short Film about Building the Simons Observatory (Part 1)




October 03, 2017


The Eternal Sky, Part 2: The Hunt for Inflation

CMB-S4 endorsed by Astro2020





Large Aperture Telescope (LAT)



Detection of **r** would give the energy scale of inflation, provide evidence for the quantization of gravity, and fundamental insights into physics and cosmology.

All inflation models that naturally explain the observed deviation from scale invariance and that also have a characteristic scale equal to or larger than the gravitational mass scale predict $r > 10^{-3}$. A well-motivated sub-class within this set of models is detectable by CMB-S4 at 5σ .

CMB-S4 sensitivity of $\sigma(r)$ < 0.0005 and ensures that a non-detection of r will rule out the leading inflationary models, and motivate alternate models for the origin of the universe.

CMB-S4 upper limit goal $r < 10^{-3}$ at 95% C.L. (SPO and SO goals ~10⁻²)

Questions?



Feel free to email me or ask questions in our eCampus Forum